

# Letters to the Editor

Innov Clin Neurosci. 2017;14(9–10):3

## AN OPEN-LABEL PROPHYLAXIS STUDY OF LITHIUM PLUS EXTENDED-RELEASE CARBAMAZEPINE (ERC-CBZ) COMBINATION FOR RAPID CYCLING BIPOLAR DISORDER

Dear Editor:

Rapid cycling bipolar disorder involves the occurrence of four or more episodes of mania and/or major depression within a year and is characterized by a treatment-refractory course with high morbidity. As rapid cycling patients often do not respond well to lithium monotherapy, mood stabilizer combinations have been recommended for long-term treatment.<sup>1</sup> Positive findings have been reported in rapid cycling patients treated with lithium combined with mood stabilizers such as carbamazepine as compared to monotherapy.<sup>2,3</sup> Extended-release carbamazepine (ERC-CBZ) has important advantages over immediate-release carbamazepine, including reduced fluctuations in serum carbamazepine levels and a decrease in CNS side effects.<sup>4</sup> Our objective was to evaluate the safety and efficacy of lithium plus ERC-CBZ for prophylaxis in patients with rapid cycling bipolar disorder.

We recruited 16 patients with DSM-IV defined bipolar disorder with a history of rapid cycling in the past 12 months and on stable lithium therapy for six months or longer. In the preliminary study phase, patients received ERC-CBZ at starting doses ranging from 100mg to 200mg twice daily depending on clinical presentation and titrated up to a maximum dose of 1,600mg/day over a two-week period. All psychotropics other than lithium, ERC-CBZ, and benzodiazepines were then tapered over a two-week period. Patients stabilized on lithium and ERC-CBZ then entered a six-month open-label phase and completed biweekly visits. The primary efficacy measure was time to relapse as estimated

using the Kaplan-Meier method. Relapse was defined by (a) a need for additional pharmacotherapy for affective symptoms, (b) hospitalization for an affective episode, and (c) increase of more than 50 percent in Hamilton Depression Rating Scale (HAM-D) and Young Mania Rating Scale (YMRS) scores from baseline. Secondary efficacy measures included changes in symptom severity on the HAM-D, YMRS, and Clinical Global Impression (CGI) scales.

Of the 16 patients who enrolled in the study, 56 percent were men ranging in age from 28 to 59 years ( $M=39.4$ ). At baseline, the mean HAM-D score was 10.6 ( $SE=1.7$ ), and the mean YMRS score was 9.6 ( $SE=2.4$ ). Twelve patients entered the open-label phase. Six patients experienced a relapse with the median time to relapse estimated as 144 days. One patient was lost to follow-up at 14 days, one patient dropped out of the study for SAE at 61 days, and four patients completed the study. The 100-day relapse rate was 27 percent (95% CI: 10–63%). Using repeated measures and a mixed-effects model, HAM-D total score, YMRS total score, and CGI severity and improvement scores did not change significantly between visits 6 and 18 ( $p>0.1$ ).

Rapid cycling bipolar disorder is highly resistant to treatment, although research evidence regarding optimal treatment for rapid cycling patients has been limited.<sup>5</sup> The findings from this study suggest that lithium combined with ERC-CBZ might be safe and effective for prophylaxis in patients with rapid cycling bipolar disorder, although the small sample size and high lost-to-follow-up rate are clear limitations. Randomized controlled trials are needed to validate these preliminary findings.

## REFERENCES

1. Calabrese JR, Shelton MD, Rapport DJ, et al. Current research on rapid cycling

- bipolar disorder and its treatment. *J Affect Disord.* 2001;67(1-3):241–255.
2. Denicoff KD, Smith-Jackson EE, Disney ER, et al. Comparative prophylactic efficacy of lithium, carbamazepine, and the combination in bipolar disorder. *J Clin Psychiatry.* 1997;58(11):470–478.
3. Di Costanzo E, Schifano F. Lithium alone or in combination with carbamazepine for the treatment of rapid-cycling bipolar affective disorder. *Acta Psychiatr Scand.* 1991;83(6):456–459.
4. Miller AD, Krauss GL, Hamzeh FM. Improved CNS tolerability following conversion from immediate- to extended-release carbamazepine. *Acta Neurol Scand.* 2004;109(6):374–377.
5. Fountoulakis KN, Kontis D, Gonda X, Yatham LN. A systematic review of the evidence on the treatment of rapid cycling bipolar disorder. *Bipolar Disord.* 2013;15(2):115–137.

With regard,

**Sriram Ramaswamy, MD, David Driscoll, PhD, and Lynette M. Smith, PhD**

Dr. Ramaswamy is with the Department of Psychiatry, Creighton University School of Medicine, Omaha, Nebraska, and the VA Nebraska-Western Iowa Health Care System, Omaha, Nebraska. Dr. Driscoll is with the VA Nebraska-Western Iowa Health Care System, Omaha, Nebraska. Dr. Smith is with the College of Public Health, University of Nebraska Medical Center, Omaha, Nebraska.

**Address for correspondence:** Sriram Ramaswamy, MD, Department of Psychiatry, Creighton University School of Medicine, 3528 Dodge Street, Omaha, NE 68131; Phone: 402-345-7100; Fax: 402-345-8815; Email: SriramRamaswamy@creighton.edu

**Keywords:** carbamazepine, lithium, rapid cycling, bipolar disorder

**Funding/financial disclosures:** This work was supported by an investigator-initiated grant from Shire Pharmaceuticals. **ICNS**